

Circadian Variation - Increased Morning Incidence of Acute Myocardial Infarction in Patients with Coronary Artery Disease

M. S. Khan, S. I. Ahmad*

Final Year Medical Student, Department of Cardiology*, Ziauddin Medical University, Karachi.

FULL TEXT

Introduction

Despite recent encouraging declines in the mortality rate in both developed and developing countries, Coronary Artery Disease (CAD) remains a preeminent cause of death and disability. While numerous modifiable and non-modifiable risk factors have been associated with an increase risk of coronary disease. Many studies in the past decade have demonstrated diurnal variation in the onset of acute cardiovascular disorders, such as myocardial ischemia¹⁻³, acute myocardial infarction (AMI)⁴⁻⁸ cardiac arrhythmias^{9,10} and sudden cardiac death.¹¹⁻¹³ These results consistently support the view that there is an increased incidence of acute cardiac events in the morning hours and a low incidence at nights as judged on the basis of cardiac symptoms. The present study also examines that a significant circadian variation exist in the onset of symptoms of Acute Myocardial Infarction in patients with coronary artery disease (CAD). The hypothesis of this study was that more than 50% of the patients with AMI present in the morning hours.

Method

Study Population

This study was based on onset of clinical symptoms and admission to the hospital. The study population included patients admitted to the coronary care units of two teaching hospitals in the city of Karachi, Dr. Ziauddin Medical University Hospital and National Institute of Cardiovascular Disease. The time of the onset was assessed by a semi-structured questionnaire which included specific question on the precise time of onset of the clinical symptoms. Questionnaire was pre-tested and all the participants gave their informed oral consent voluntarily, this criterion excluded the patients from Emergency Department because of their condition. All the consecutive patients who were diagnosed as having acute myocardial infarction and willing to participate, were included in the study. The diagnosis of AMI was made if at least two of the following three predefined criteria were satisfied: a history of chest pain consistent of myocardial ischemia; enzymes elevations of creatinine kinase, its isoenzyme subfraction or lactate dehydrogenase or positively Troponin- T tests and in ECG tracings showing ischemic changes in the "ST" segment or presence of any new "Q" wave, typical of AMI.

Statistical Analysis

The computer package "Epi-Info version 6" was used for data entry and analysis. To determine whether or not the incidence of myocardial infarction exhibited a systematic circadian variation, the following statistical approaches were prospectively chosen. Test of proportion was used to test the difference between the incidence of AMI in two groups of morning and evening. Statistically significant difference was defined as P value of less than 0.05.

Results

Sixty-nine men and thirty-one women with confirmed AMI were enrolled in the study. The mean age was 55 years \pm 15. The incidence of Myocardial infarction assessed by the onset of clinical symptoms and the time of admission exhibited a marked circadian variation with a peak period during morning hours. (Figure). Myocardial infarction occurred more frequently (74% vs. 26%, $p < 0.001$) (Table 1) during morning (period of maximum incidence) hours in the early morning hours as compared to evening hours

(period of minimum incidence). There were also very few cases of AMI, reported in the subsequent hours but they do not approximate the increased frequency observed during morning hours. The circadian pattern in the incidence of Myocardial infarction was similar among older and young patients and in male and female patients. (Table 1). Troponin T was found to be the most accurate diagnostic test, detecting about 89% of the cases of acute myocardial infarction. Males had more AMI compared to females. Hypertension was found to be the most common risk factor (47%)(Table 2). The most common presenting symptoms were retrosternal chest pain (91%) followed by dyspnea and others (Table 3). The Myocardial walls, which were more vulnerable to get infarct, are anterior wall (76%) followed by inferior (22%) and posterior walls (2%).

Discussion

The results of this study suggest an increased incidence of the onset of initial symptoms of AMI in morning hours. (Figure). This morning peak was shown consistently in different subgroups, which indicated that it is a phenomenon not attributable to risk variables alone. There was no indication of further periodicity except for the morning peak. Although a circadian variation in the onset of Myocardial Infarction is pointed out in several previous studies¹⁴⁻¹⁶, this study also extends previous observation. The factor responsible for this observation remains elusive for the most part and warrant further investigations. However, there are several clues to the factors responsible for this phenomenon. Platelets aggregation has been reported to be more responsive during morning.¹⁷⁻¹⁸ Conceivably, early morning hypercoagulability could enhance the intracoronary thrombosis and there by increases the risk of major advance cardiac events (MACE).¹⁹⁻²¹ Another important factor in the pathogenesis of the increased morning incidence of myocardial infarction may be the early morning increase of arterial pressure.²²⁻²³ The increased arterial blood pressure, increases the myocardial oxygen demand that leads to the increased vulnerability of the myocardium to ischemia, secondly the increase in arterial blood pressure could enhance the risk of plaque fissure that has been shown to occur rapidly before myocardial infarction.²⁴ Several other factors for e.g. hormonal changes²⁵ or increase in coronary artery tone^{26,27} may also play a role in the pathogenesis of this phenomenon. It has been proposed that in a number of cases, sudden cardiac death is the result of primary arrhythmic event.²⁸ Such fatal arrhythmias are more likely to occur in the morning since increased activity of the sympathetic nervous system at that time.²⁹ which in turn may increase electrical instability and subsequent conduction defects.³⁰ It has recently been suggested that there is a parallel reduction in incidence of myocardial infarction and sudden cardiac death in patients on aspirin prophylaxis or beta blocker therapy.^{31,32}

The limitation of the present investigation includes its small sample size, lack of enzymatic determination of the onset of AMI, failure to collect information from patients dying soon after hospital admission or too ill to provide informed consent. However this study does provide sufficient baseline data on which further work can be based.

Conclusion

Quiet obvious enough, morning appears to be a time when the "transient risk factors" that lead to sudden cardiac phenomenon are likely to be prominent. The present findings help to narrow the search for the possible cause or causes of the circadian variation in the onset of AMI. It is necessary to further investigate the physiological changes involved to eventually design more effective and preventive therapy and to reduce the incidence of the disorder.

The recommendation that should be emphasized are: Anti platelet therapy reduces early morning myocardial infarction³¹, keeping the importance of platelet characteristics in mind. Prevention of blood pressure surge in the morning and limitation of myocardial oxygen demand may be protected by giving beta adrenergic blocking therapy ³² and long acting anti ischemic medication perhaps if taken at night provides better protection. From this study it is obvious that, the potential triggering activities such as extreme physical exertion, mental stress and early morning sexual activity should be avoided.^{33,34} Para-medical staff should be trained accordingly to manage early morning cardiac emergencies.

Acknowledgements

This study was made possible by the cooperation of the cardiology staffs of the two hospitals; Dr. Ziauddin Medical University Hospital and National Institute of Cardiovascular Disease. This study would never be completed without the sincere efforts and prayers of my family and friends.

References

1. Portman M. A. Molecular clock mechanisms and circadian rhythms intrinsic to the heart. *Circ Res* 2001;89:1084-86.
2. Mickley H, Pless P, Nielsen JR, et al. Circadian variation of transient myocardial ischemia in the early out-of-hospital period after first acute myocardial infarction. *Am J Cardiol* 1991;67:927-32.
3. Andrews TC, Fenton T, Toyosaki N, et al. Subsets of ambulatory myocardial ischemia based on heart rate activity: circadian distribution and response to anti-ischemic medication. *Circulation* 1993;88:92100.
4. Goldberg RJ, Brady P, Muller JE, et al. Time of onset of symptoms of acute myocardial infarction. *Am J Cardiol* 1990;66:140-4.
5. Ridker PM, Manson JE, Buring JE, et al. Circadian variation of acute myocardial infarction and the effect of low-dose aspirin in a randomized trial of physicians. *Circulation* 1990;82:897-902.
6. Willich SN, Lowel H, Lewis M, et al. Association of wake time and the onset of myocardial infarction: triggers and mechanisms of myocardial infarction (TRIMM) pilot study. *Circulation* 1991; 84:62-7.
7. Hansen O, Johansson BW, Gullberg B. Circadian distribution of onset of acute myocardial infarction in subgroups from analysis of 10,791 patients treated in a single center. *Am J Cardio*. 1992;69:1003-8.
8. Tofler GH, Muller JE, Stone PH, et al. Modifiers of timing and possible triggers of acute myocardial infarction in the thrombolysis in myocardial infarction phase II (TIMI II) study group. *J Am Coll Cardiol* 1992;20:1049-55.
9. Siegel D, Black DM, Seeley DG, et al. Circadian variation in ventricular arrhythmias in hypertensive men. *Am J Cardiol* 1992;69:344-7.
10. Lampert R, Rosenfeld L, Batsford W, et al. Circadian variation of sustained ventricular tachycardia in patients with coronary artery disease and implantable cardioverter-defibrillators. *Circulation* 1994;90:241-7.
11. Arntz HR, Willich SN, Oeff M, et al. Circadian variation of sudden cardiac death reflects age-related variability in ventricular fibrillation. *Circulation* 1993;88: 2284-9.
12. Aronow WS. Circadian variation of primary cardiac arrest or sudden cardiac death in patients aged 62 to 100 years (mean 82). *Am J Cardiol* 1993;71:1455-6.
13. Maron BJ, Kogan J, Proschan MA, et al. Circadian variability in the occurrence of sudden cardiac death in patients with hypertrophic cardiomyopathy. *J Am Coll Cardio*. 1994;23:1405-9.
14. Robert JG, Priscilla B, James EM. Time of onset of symptoms of acute myocardial infarction. *Am J Cardiol* 1990;66:140-4.
15. Muller JE, Ludmer PL, Willich SN, et al. Circadian variation in the frequency of sudden cardiac death. *Circulation* 1987; 75:131-8.
16. Rocco MB, Barry J, Campbell S, et al. Circadian variation of transient myocardial ischemia in patients with coronary artery disease. *Circulation* 1987;75:395-400.

17. Kapiotis S, Jilma B, Quehenberger P, et al. Morning Hypercoagulability and Hypofibrinolysis: Diurnal Variations in Circulating Activated Factor VII, Prothrombin Fragment F1+2, and Plasmin-Plasmin Inhibitor Complex. *Circulation* 1997;96: 19-21.
18. Siess W, Lorenz R, Roth P, et al. Plasma catecholamines, platelet aggregation and associated thromboxane formation after physical exercise, smoking or norepinephrine infusion. *Circulation* 1982 66:44-8.
19. Ehrly AM, Jung G. Circadian rhythm of human blood viscosity. *Biorheology* 1973;10:577-83.
20. Kubota K, Sakurai T, Tamura J, et al. Is the circadian change in hematocrit and blood viscosity a factor triggering cerebral and myocardial infarction? *Stroke* 1987;18:812-13.
21. Tofler GH, Brezinski D, Schafer AI, et al. Morning increase in platelet response to ADP and epinephrine: association with the time of increased risk of myocardial infarction sudden cardiac death. *N Engl J Med* 1987;316:1514 -18.
22. Millar-Craig MW, Bishop CN, Reftery EB: Circadian variation of blood pressure. *Lancet* 1978;1:795-7.
23. Meyer-Sabellek W, Schulte KL, Distler A, et al. Followup of a method of twenty-four-hour indirect blood pressure monitoring. *Nephron* 1987;47:42-6.
24. Davies MJ, Thomas AC. Plaque fissuring; The case of acute myocardial infarction, sudden ischemic death and crescendo angina. *Br Heart J* 1985;53:363-73.
25. Dodt C, Breckling U, Derad, et al. Plasma Epinephrine and Norepinephrine Concentrations of healthy humans associated with nighttime sleep and morning arousal. *Hypertension* 1997; 30:71-6.
26. Yeung AC, Vekshtein VI, Krantz DS, et al. The effect of atherosclerosis on the vasomotor response of coronary arteries to mental stress. *N Engl J Med* 1991;325:1551-6.
27. El-Tamimi H, Mansour M, Pepine C, et al. Circadian variation in coronary tone in patients with stable angina : protective role of the endothelium. *Circulation* 1995;92: 3201-5.
28. Peters RW, Zoble RG, Liebson PR, et al. Identification of a secondary peak in myocardial infarction onset 11 to 12 hours after awakening: the cardiac arrhythmia suppression trial (CAST) experience. *J Am Coll Cardiol.* 1993; 22:998-1003.
29. Muller JE, Tofler GH, Verrier RL. Sympathetic activity as the cause of the morning increase in cardiac events: a likely culprit, but the evidence remains circumstantial. *Circulation* 1995;91:2508-9.
30. Kupari M, Koskinen P, Leinonen H. Double-peaking circadian variation in the occurrence of sustained supraventricular tachyarrhythmias. *Am Heart J* 1990;120:1364-9.
31. Mulcahy D, Keegan J, Cunnigham J, et al. Circadian variation of total ischemic burden and its alteration with anti-anginal agents. *Lancet* 1988;2:755-9.
32. Willich SN, Linderer T, Wegscheider K, et al. ISAM Study Group. Increased morning incidence of myocardial infarction in the ISAM study: absence with prior β -adrenergic blockade. *Circulation* 1989;80:853-8.
33. Willich SN, Lewis M, Lowel H, et al. The triggers and mechanisms of myocardial infarction. Physical Exertion as a trigger of acute myocardial infarction. *N Engl J Med* 1993;329:1684-90.
34. Rozanski A, Bairey CN, Krantz DS, et al. Mental stress and the induction of silent myocardial ischemia in patients with coronary artery disease. *N Engl J Med* 1988; 318:1005-12.